

A review of actin binding proteins: new perspectives

Ricardo Uribe · David Jay

Received: 29 June 2007 / Accepted: 28 September 2007 / Published online: 16 October 2007
© Springer Science+Business Media B.V. 2007

Abstract Actin binding proteins (ABPs) have been considered components of the cytoskeleton, which gives structure and allows mobility of the cell. The complex dynamic properties of the actin cytoskeleton are regulated at multiple levels by a variety of proteins that control actin polymerization, severing of actin filaments and cross-linking of actin filaments into networks, which may be used by molecular motors. Proteins that cross-link F-actin are important for the maintenance of the viscoelastic properties of the cytoplasm and for the integrity of plasma membrane-associated macromolecules. Most of these F-actin cross-linking proteins have an actin-binding domain homologous to calponin. In addition, some of them have been considered scaffolds. Through the years, several research groups have found different proteins that interact with ABPs; however, the effect of these interactions on ABPs remains mostly unknown. In addition to organize the cytoskeletal structure, recent data indicate that ABPs can also migrate to the nucleus. This fact is in agreement and could be relevant to the recently found role that actin might play in nuclear function. Recent data and analysis of published results have also indicated that scaffold proteins like filamin A (FLNa) may be processed by proteolysis and that the degradation products generated by this reaction may play a role as signaling molecules, integrating nuclear and cytosolic pathways. Some of the relevant information in this area is reviewed here.

Keywords Actin · Actin binding proteins (ABPs) · Calpain · Nucleus · Proteolysis · Signaling integrator proteolytic peptides (SIPPs)

Abbreviations

ABPs	Actin binding proteins
ABD	Actin-binding domain
CH domains	Calponin homology domains
FLNa	Filamin A
SIPPs	Signaling integrator proteolytic peptides

Actin in the nucleus

The actin cytoskeleton of eukaryotic cells plays a central role in many cell functions such as cell shape, cell division, motility, contraction, adhesion, phagocytosis, protein sorting and signal transduction. The complex and dynamic properties of the actin cytoskeleton are regulated at multiple levels by a variety of proteins that control actin polymerization, severing of actin filaments and cross-linking of actin filaments into networks or bundles. Apart from being a fundamental component of cytoskeleton, actin has been found in the cellular nucleus carrying out different functions. Many studies have established the presence of actin in the nucleus and have shown that its functions are as diverse here as the ones found in the cytoplasm. Possible roles for nuclear actin include contribution to the organization of chromatin remodeling complexes, RNA processing, or regulation of DNase I function [1, 2]. In addition, actin plays a direct role in transcription by RNA polymerases I, II and III [3–6]. Until now the mechanism by which actin migrates to the nucleus is unknown. In this

R. Uribe (✉) · D. Jay
Departamento de Biomedicina Molecular Cardiovascular,
Instituto Nacional de Cardiología Ignacio Chávez, Juan Badiano
No. 1 Colonia Seccion XVI, Tlalpan, 14080 Mexico, DF,
Mexico
e-mail: rickurib@yahoo.com.mx

regard, myosin has been found with actin in the nucleus, physically associated with both RNA polymerase I holo-enzyme and the ribosomal genes [3]. The ability of myosin to bind actin is shared by other actin binding proteins (ABPs), that contain actin-binding domains (ABD), and these ABPs may localized to the nucleus too. Filamentous actin (F-actin) cross-linker proteins are very abundant in normal cells. The possibility exists, therefore, that these proteins migrate to the nucleus together with actin by means of these ABDs and in this way they may transport actin to the nucleus. In this regard, however, it is worthwhile mentioning that there is a growing body of evidence indicating that nuclear actin is not in a conventional form but rather it is present as short oligomers. In this sense, ABPs would not provide a nucleating unit around which an F-actin network could be organized; especially in the cases where ABPs have already been digested (see below). Therefore, nuclear actin, alone or in complex with ABPs, would provide a function different from the structural one found in the cytoplasm [7, 8].

Actin binding proteins with CH domains

There is a great diversity of proteins that bind actin, which are important contributors to cell structure. These proteins are important in maintaining the viscoelastic properties of the cytoplasm, to protect the integrity of the plasmatic membrane-associated macromolecules and to serve as scaffolds. In eukaryotic cells, there are diverse F-actin cross-linking proteins. Many of these proteins have actin binding domains composed of two calponin homology

(CH) regions. The CH domain is a protein domain of ~ 110 residues found in cytoskeletal and signal-transduction proteins [9]. Two CH domains in tandem constitute the actin-binding region of members of several protein families that cross-link actin filaments forming networks or bundles. These families include proteins such as spectrin, α -actinin, dystrophin and FLNa (Fig. 1). The surface of the repeats of these proteins may serve as docking sites for cytoskeletal and signal transduction proteins. For a long time the idea existed that these ABPs were exclusively found associated to the cytoskeleton in the cytoplasm, but evidence has arisen recently that demonstrates that they are also found in the nucleus carrying out diverse functions.

ABPs in the nucleus: a reality

The demonstration of the location of a protein that binds actin in the nucleus was published for the first time in 1987 [10]. Thereafter, other actin-binding proteins have been located in the nucleus (Table 1).

α -actinin interacts with more than 29 components in the cell [11], and there are different isoforms of this protein, like α -actinin-4, that cross-links actin filaments and reorganizes the cytoskeleton for cell movement. Nevertheless, it has also been observed that α -actinin-4 is present in the nucleus of a particular population of breast cancer cells. However, from the deduced aminoacid sequence, it seems that actinin-4 does not possess any apparent nuclear localization signal [12]. Likewise, Pinaev's research group found that the α -actinin-4 co-localizes to the nucleus with p65/RelA subunit of NF-kappa B [13]. In addition,

Fig. 1 Schematic representation of the different proteins that bind actin. Some similarities in the structure can be appreciated. These proteins are formed by repetitions, (alpha or beta) of approximately 100 aminoacids. The CH domains are present in most of these proteins




Table 1 Actin binding proteins with CH domains found in the nucleus

Actin binding proteins	References
Alpha actinin-4	[12–14]
α -Spectrin II	[16–18, 20]
β -spectrin II	[19]
Filamin A	[23, 24, 27]
Dystrophin	[31]
Vav	[32]
Nesprin-2	[35]
L-Plastin	[36]

α -actinin 4 potentiates myocyte enhancer factor-2 transcription activity by antagonizing histone deacetylase 7 [14]. These results demonstrate that this ‘cytosol-associated cytoskeletal protein’ has important functions in the nucleus.

Another cytoskeletal F-actin cross-linking protein localized to the nucleus is spectrin. This protein has diverse isoforms. Spectrin was also detected at the nuclear envelope, as well as in the intranuclear granules within liver cells [15].

Strong evidence exists indicating that α -SpecII isoform of spectrin plays a role in DNA repair, possibly acting as a scaffold for DNA repair proteins [16–18]. Other isoform of spectrin, β -SpecII, is present in the nucleus and interact with nuclear proteins. β -SpecII associates with Smad proteins in the cytoplasm and once associated, these proteins can translocate to the nucleus of liver cells [19].

Deepa M. Sridharan et al. have observed that nonerythroid α -spectrin II Σ^* (α SpII Σ^*) associates with five different groups of functionally important proteins in the nucleus of normal human cells: structural proteins, proteins involved in DNA repair, chromatin remodeling proteins, Fanconi anemia proteins and proteins which play a role in transcription and processing of RNA [20].

Another cytoskeletal-associated protein that does not belong to the spectrin family, but has also been located to the nucleus is filamin A (FLNa). Filamins are homodimeric F-actin cross-linkers, with a high molecular weight (280 kDa for FLNa), which organize actin filaments into parallel arrays or three-dimensional webs, linking them to cellular membrane [21]. This protein anchors a variety of transmembrane proteins to the actin cytoskeleton and provides a scaffold for many cytoplasmic and signaling molecules [22].

Ozanne and Loy observed that the C-terminal region of the FLNa could translocate to the nucleus together with the androgen receptor (AR) [23, 24]. The C-terminal region of FLNa is a 100-kDa polypeptide. This polypeptide can be generated by the action of calpain when FLNa is dephosphorylated in Ser 2152 [25, 26]. FLNa can migrate with other

proteins to the nucleus, for example Protein FOXC1 [27]. Although in this case, it remains to be established whether full length FLNa or a proteolysis fragment of it is the migrating element. Furthermore, FLNa regulates the subcellular localization of Smad2 and PEBP2/CBF, two transcription factors that are usually found in the cytoplasm [28, 29].

There is a growing body of evidence implicating FLNa in the regulation of nuclear functions including DNA repair [30].

Another ABP with a CH domain is Dystrophin 71 (Dp71), which was reported to be found in the nucleus of PC12, cells an established cell line derived from a rat pheochromocytoma [31].

At this point the question arises as to whether all these ABPs could migrate to the nucleus associated with actin. In this regard, it is worthwhile mentioning that FLNa was localized in the nucleus by Fred Berry et al. utilizing an antibody against the N-terminal actin-binding region of the protein [27].

It is now clear that ABPs and actin perform a variety of functions in the nucleus. How these proteins are mobilized between the nucleus and the cytoplasm and whether they control nuclear processes on their own or in association with each other, remains as open questions. Although there is no final evidence indicating that an association of this kind can play a part in the mobilization of actin to the nucleus, this possibility cannot be ruled out at the moment since different ABPs with CH domain have been found in the nucleus. For example, *vav*, a 95 kDa CH-containing GDP/GTP exchange proto-oncogene product expressed exclusively in hematopoietic cells was found to bind to Heterogeneous Nuclear Ribonucleoprotein (hnRNP) C through a region in the C-terminal domain of the protein. The interaction was regulated by poly(U) RNA and was proposed to play a part in pre-mRNA synthesis and mRNA export [32]. Whether *vav* was bound to hnRNP C alone or in complex with actin through its CH domain remains as an open question. However, it is worthwhile mentioning that more recent results showed that actin and hnRNP U cooperate for productive transcription by RNA polymerase II [33]. In this case hnRNP U also bound actin through a short amino acid sequence in the C-terminal region. In addition, Laury-Kleintop et al. recently reported that hnRNP K form a signaling complex with calponin that may be involved in growth-stimulated post-transcriptional regulation. In this case the complex was found mainly in a cytoplasmic location [34].

How can the ABPs arrive at the nucleus?

Since it was known that ABPs could be localized to the nucleus, the search started for a possible mechanism for

their mobilization. There has not been final evidence indicating whether full-length ABPs are mobilized within the cell and to the nucleus or only fragments of them, probably produced by protease action, are required to fulfill the function. Previous studies have demonstrated that in this case the susceptibility of some of ABPs can be regulated by phosphorylation as in the case of the FLNa or alpha II spectrin [25, 26, 37]. In the case of FLNa, two major peptides are generated after proteolysis, corresponding to the C-terminal 100 kDa and N-terminal 180 kDa regions of the molecule. In this regard, Ozanne and Loy found that the C-terminal region of FLNa could be translocated to the nucleus associated to the AR [23, 24]. Previous studies demonstrated that the generation of this C-terminal fragment of FLNa could be prevented by the phosphorylation of Ser 2152 by PKA [25, 26].

The results of Ozanne and Loy indicated that at least one region of the protein could carry out a specific function, i.e. mobilization of the AR to the nucleus. Most of the ABPs are also susceptible to degradation by caspases and could be also the target or other proteases.

We have recently proposed that proteolysis of scaffold proteins and other possible targets could provide a general mechanism to integrate cellular pathway within the cell. The protein fragments produced by proteolysis could function as “Signaling Integrator Proteolytic Peptide” (SIPPs) (Fig. 2), and their generation could be regulated by a phosphorylation/dephosphorylation processes [38]. We anticipate that many of the ABPs could also participate in

nuclear and cytoplasmic routs by these means. Thus, it is possible that ABP SIPPs alone or in combination with actin could serve as nuclear scaffolds to proteins involved in DNA repair or chromatin complex formation.

The study of the nuclear ABPs could probably require the analysis of the proteolytic fragments generated from them that could represent the true functional peptides.

References

1. Olave IA, Reck-Peterson SL, Crabtree GR (2002) Nuclear actin and actin-related proteins in chromatin remodeling. *Annu Rev Biochem* 71:755–781
2. Chhabra D, Bao S, dos Remedios CG (2002) The distribution of cofilin and DNase I in vivo. *Cell Res* 12(3–4):207–214
3. Fomproix N, Percipalle P (2004) An actin-myosin complex on actively transcribing genes. *Exp Cell Res* 294(1):140–148
4. Philimonenko VV, Zhao J, Iben S, Dingova H, Kysela K, Kahle M, Zentgraf H, Hofmann WA, de Lanerolle P, Hozak P, Grummt I (2004) Nuclear actin and myosin I are required for RNA polymerase I transcription. *Nat Cell Biol* 6(12):1165–1172
5. Hofmann WA, Stojiljkovic L, Fuchsova B, Vargas GM, Mavrommatis E, Philimonenko V, Kysela K, Goodrich JA, Lessard JL, Hope TJ, Hozak P, de Lanerolle P (2004) Actin is part of pre-initiation complexes and is necessary for transcription by RNA polymerase II. *Nat Cell Biol* 6(11):1094–1101
6. Hu P, Wu S, Hernandez N (2004) A role for beta-actin in RNA polymerase III transcription. *Genes Dev* 18:3010–3015
7. Jockusch BM, Schoenenberger C-A, Stetefeld J, Aebi U (2006) Tracking down the different forms of nuclear actin. *Trends Cell Biol* 16(8):391–396
8. Schoenenberger CA, Buchmeier S, Boerries M, Sütterlin R, Aebi U, Jockusch BM (2005) Conformation-specific antibodies reveal distinct actin structures in the nucleus and the cytoplasm. *J Struct Biol* 152(3):157–168
9. Stradal T, Kranewitter W, Winder SJ, Gimona M (1998) CH domains revisited. *FEBS Lett* 431(2):134–137
10. Nishida E, Iida K, Yonezawa N, Koyasu S, Yahara I, Sakai H (1987) Cofilin is a component of intranuclear and cytoplasmic actin rods induced in cultured cells. *Proc Natl Acad Sci USA* 84:5262–5266
11. Otey CA, Carpen O (2004) Alpha-actinin revisited: a fresh look at an old player. *Cell Motil Cytoskeleton* 58(2):104–111
12. Honda K, Yamada T, Endo R, Ino Y, Gotoh M, Tsuda H, Yamada Y, Chiba H, Hirohashi S (1998) Actinin-4, a novel actin-bundling protein associated with cell motility and cancer invasion. *J Cell Biol* 140(6):1383–1393
13. Babakov VN, Bobkov DE, Petukhova OA, Kropacheva IV, Podol'skaia EP, Pinaev GP (2004) Alpha-Actinin-4 and p65/RelA subunit of NF-kappaB transcription factor are co-localized and migrate together into the nucleus in EGF-stimulated A431 cell. *Tsitologiya* 46(12):1064–1072
14. Chakraborty S, Reineke EL, Lam M, Li X, Liu Y, Gao C, Khurana S, Kao HY (2006) Alpha-actinin 4 potentiates myocyte enhancer factor-2 transcription activity by antagonizing histone deacetylase 7. *J Biol Chem* 281(46):35070–35080
15. Bachs O, Lanini L, Serratosa J, Coll MJ, Bastos R et al (1990) Calmodulin-binding proteins in the nuclei of quiescent and proliferatively activated rat liver cells. *J Biol Chem* 265: 18595–18600
16. McMahon LW, Walsh CE, Lambert MW (1999) Human alpha spectrin II and the Fanconi anemia proteins FANCA and FANCC




Fig. 2 Schematic representation of SIPPs generation. SIPPs from ABPs or other targets can translocate to the nucleus together with different proteins

- interact to form a nuclear complex. *J Biol Chem* 274: 32904–32908
17. McMahon LW, Sangerman J, Goodman SR, Kumaresan K, Lambert MW (2001) Human alpha spectrin II and the FANCA, FANCC, and FANCG proteins bind to DNA containing psoralen interstrand cross-links. *Biochemistry* 40:7025–7034
 18. Sridharan D, Brown M, Lambert WC, McMahon LW, Lambert MW (2003) Nonerythroid alphaII spectrin is required for recruitment of FANCA and XPF to nuclear foci induced by DNA interstrand cross-links. *J Cell Sci* 116:823–835
 19. Tang Y, Katuri V, Dillner A, Mishra B, Deng CX (2003) Disruption of transforming growth factor-beta signaling in ELF beta-spectrin-deficient mice. *Science* 299:574–577
 20. Sridharan DM, McMahon LW, Lambert MW (2006) alphaII-Spectrin interacts with five groups of functionally important proteins in the nucleus. *Cell Biol Int* 30(11):866–878
 21. Gorlin JB, Yamin R, Egan S, Stewart M, Stossel TP, Kwiatkowski DJ, Hartwig JH (1990) Human endothelial actin-binding protein (ABP-280, nonmuscle filamin): a molecular leaf spring. *J Cell Biol* 111:1089–1105
 22. Pudas R (2006) Structural and interaction studies on the carboxy-terminus of filamin, an actin binding protein. Dissertation, University of Oulu
 23. Ozanne DM, Brady ME, Cook S, Gaughan L, Neal DE, Robson CN (2000) Androgen receptor nuclear translocation is facilitated by the f-actin cross-linking protein filamin. *Mol Endocrinol* 14(10):1618–1626
 24. Loy CJ, Sim KS, Yong EL (2003) Filamin-A fragment localizes to the nucleus to regulate androgen receptor and coactivator functions. *Proc Natl Acad Sci USA* 100(8):4562–4567
 25. Jay D, Elizabeth J, Lara JE (2000) Determination of a cAMP-dependent protein kinase phosphorylation site in the C-terminal region of human endothelial actin-binding protein. *Arch Biochem Biophys* 377(1):80–84
 26. Jay D, Elizabeth J (2004) In situ determination of a PKA phosphorylation site in the C-terminal region of Filamin. *Mol Cell Biochem* 260:49–53
 27. Berry FB, O'Neill MA, Coca-Prados M, Walter MA (2005) FOXO1 transcriptional regulatory activity is impaired by PBX1 in a Filamin A-mediated manner. *Mol Cell Biol* 25(4):1415–1424
 28. Sasaki A, Masuda Y, Ohta Y, Ikeda K, Watanabe K (2001) Filamin associates with Smads and regulates transforming growth factor-beta signaling. *J Biol Chem* 276:17871–17877
 29. Yoshida N, Ogata T, Tanabe K, Li S, Nakazato M, Kohu K, Takafuta T, Shapiro S, Ohta Y, Satake M, Watanabe T (2005) Filamin A-bound PEBP2beta/CBFbeta is retained in the cytoplasm and prevented from functioning as a partner of the Runx1 transcription factor. *Mol Cell Biol* 25(3):1003–1012
 30. Meng X, Yuan Y, Maestas A, Shen Z (2004) Recovery from DNA damage induced G2 arrest requires the actin-binding protein Filamin-A/actin-binding protein 280. *J Biol Chem* 279: 6098–6105
 31. Fuentes-Mera L, Rodriguez-Munoz R, Gonzalez-Ramirez R, Garcia-Sierra F, Gonzalez E, Mornet D, Cisneros B (2006) Characterization of a novel Dp71 dystrophin-associated protein complex (DAPC) present in the nucleus of HeLa cells: members of the nuclear DAPC associate with the nuclear matrix. *Exp Cell Res* 312(16):3023–3035
 32. Romero F, Germani A, Puvion E, Camonis J, Varin-Blank N, Gisselbrecht S, Fischer S (1998) Vav Binding to Heterogeneous Nuclear Ribonucleoprotein (hnRNP) C. *J Biol Chem* 273(10): 5923–5931
 33. Kukalev A, Nord Y, Palmberg C, Bergman T, Percipalle P (2005) Actin and hnRNP U cooperate for productive transcription by RNA polymerase II. *Nat Struct Mol Biol* 12(3):238–244
 34. Laury-Kleintop LD, Tresini M, Hammond O (2005) Compartmentalization of hnRNP-K during cell cycle progression and its interaction with calponin in the cytoplasm. *J Cell Biochem* 95(5):1042–1056
 35. Zhang Q, Ragnauth CD, Skepper JN, Worth NF, Warren DT, Roberts RG, Weissberg PL, Ellis JA, Shanahan CM (2005) Nesprin-2 is a multi-isomeric protein that binds lamin and emerin at the nuclear envelope and forms a subcellular network in skeletal muscle. *J Cell Sci* 118:673–687
 36. Delanote V, Van Impe K, De Corte V, Bruyneel E, Vetter G, Boucherie C, Mareel M, Vandekerckhove J, Friederich E, Gettemans J (2005) Molecular basis for dissimilar nuclear trafficking of the actin-bundling protein isoforms T- and L-plastin. *Traffic* 6(4):335–345
 37. Nicolas G, Fournier CM, Galand C, Malbert-Colas L, Bournier O, Kroviarski Y, Bourgeois M, Camonis JH, Dhemy D, Grandchamp B, Lecomte MC (2002) Tyrosine phosphorylation regulates alpha II spectrin cleavage by calpain. *Mol Cell Biol* 22:3527–3536
 38. Garcia E, Stracher A, Jay D (2006) Calcineurin dephosphorylates the C-terminal region of filamin in an important regulatory site: a possible mechanism for filamin mobilization and cell signaling. *Arch Biochem Biophys* 446(2):140–150